



# InjectaMax

## Multi-Mineral Injection **+ B12**

An injectable supplemental source of zinc, manganese, copper, selenium and vitamin B12 for cattle

**ACTIVE CONSTITUENTS** Each mL contains Zinc 40 mg/mL (as disodium zinc EDTA), manganese 10 mg/mL (as disodium manganese EDTA), copper 15 mg/mL (as copper disodium EDTA) plus **selenium** 5 mg/mL (as sodium selenite) and 1.4 mg/mL Vitamin B12 (cyanocobalamin)



### DIRECTIONS FOR USE

*By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.*

Dosing is by subcutaneous injection (under the skin) into the anterior half of the neck (the half of the neck closest to the head).



### READ ENTIRE CARTON BEFORE USE.

Only treat animals in good health.  
 Minimise stress to animals during handling for administration.  
 Slight local reaction may occur for about 30 seconds after injection.  
 A slight swelling may be observed at the injection site for a few days after administration.  
 Use standard aseptic procedures during administration of injections.  
 Doses greater than 7mL should be administered in separate sites.

### DOSAGE RECOMMENDATIONS

<b>Calves</b>	Up to 1 year 1mL per 50kg
<b>Cattle</b>	1-2 years 1mL per 75kg
<b>Cattle</b>	Over 2 years 1mL per 100kg
<b>Cows</b>	3 – 4 weeks before calving, mating and drying off
<b>Calves</b>	At day 1 and weaning
<b>Heifers</b>	Every 3 months, especially 3 – 4 weeks before mating
<b>Additional</b>	Every 2 months in wet conditions

### CONTRAINDICATIONS

Not to be used concurrently with any other form of copper supplementation, or administered at the same time as any other treatment e.g. drenching, vaccination. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Do not use at the same time as any other selenised fertiliser, prill, or product without consulting a veterinarian. Do not exceed the stated dose. Do not administer to emaciated animals of body condition score less than 3/10.

Available in 500mL flexi packs



**Precautions:** Not to be used concurrently with any other form of copper supplementation. Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian. Do not administer to severely debilitated animals. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Do not treat thirsty or exhausted cattle. Only treat animals in good health. Allow animals access to water after treatment.

#### **WARNINGS**

Harmful if swallowed. Causes mild skin irritation and serious eye irritation. May cause an allergic skin reaction. May cause damage to organs through prolonged or repeated exposure.

#### **HANDLING PRECAUTIONS**

Do not breathe mist. Wear protective gloves and eye/face protection. Do not eat, drink or smoke when using this product. Wash hands and exposed skin thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace.

#### **FIRST AID**

IF SWALLOWED, rinse mouth. Call the National Poisons Centre 0800 POISON (0800 764 766) or doctor immediately if you feel unwell. IF IN EYES, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice. IF ON SKIN, wash with plenty of water. If skin irritation or rash occurs, get medical advice. In the event of accidental self-injection, medical attention should be sought immediately. Take off contaminated clothing and wash it before reuse.

#### **ENVIRONMENTAL PROTECTION**

Harmful to aquatic life with long lasting effects. Hazardous to terrestrial vertebrates. Avoid release to the environment. Collect spillage.

#### **DISPOSAL**

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility.

#### **WITHHOLDING PERIODS**

No withholding period is required when used as directed.

#### **STORAGE**

Store below 25°C in original box. Keep container well closed. Once opened must be used in 1 month.

Registered pursuant to the ACVM Act 1997, No. A011926.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.  
HSNO No. HSR002521

#### **Registered to:**

**HORIZON AGRESOURCES (NZ) LTD**

Gloucester Court, 250 Gloucester St, Napier 4112.

#### **Distributed by:**

Agritrade Ltd



**Freephone 0800 333 855**

